Chirurgia (Bucur). 2013;108(5):736-740

Calcific Uremic Arteriolopathy in Hemodialyzed Patients

I.A. Checheriță[1], D. Smarandache[1], D. Rădulescu[1], I. Peride[1], O. Bratu[2], A. Ciocâlteu[1], I. Sebe[3], I. Lascăr[3]

[1]Department of Nephrology and Dialysis, Clinical Department No. 3, "Carol Davila" University of Medicine and Pharmacy Bucharest, Romania

[2]Department of Urology, Clinical Department No. 3, "Carol Davila" University of Medicine and Pharmacy Bucharest, Romania

[3]Department of Microsurgery and Plastic Surgery, Clinical Department No. 11, "Carol Davila" University of Medicine and Pharmacy Bucharest, Romania

Abstract

In the present article, we discuss the case of 67-year-old female patient diagnosed with inferior limbs calciphylaxis and hemodialyzed since 2006. The clinical manifestations and pathological lab findings are typical for this rare and extremely severe complication in chronic hemodialyzed individuals. The favorable treatment response to sodium thiosulfate, not often used as elected therapy in international studies, represents the particularity of the case.

Key words: calciphylaxis, hemodialysis, treatment

Corresponding author: Assistant Professor I.A. Checheriåã, MD, PhD
Department of Nephrology and Dialysis
Clinical Department No. 3, "Carol Davila"
University of Medicine and Pharmacy
37th Dionisie Lupu Street, 1st Sector, 020022, Bucharest, Romania
E-mail: fizij@yahoo.com