Chirurgia (Bucur). 2021; 116(1):117-122
http://dx.doi.org/10.21614/chirurgia.116.1.117

Fibroza retroperitonealã idiopaticã

Virgiliu-Mihail Prunoiu[1,2], Marian Augustin Marincaș[1,2], Eugen Brãtucu[1,2], Neicu Ștefania[3], Maria-Manuela Rãvaș[1,2]
[1]General Surgery and Oncology Clinic I, "Prof. Al. Trestioreanu" Oncology Institute, Bucharest, Romania
[2]Carol Davila University of Medicine and Pharmacy, Bucharest, Romania
[3]Department of Pathological Anatomy, "Prof. Al. Trestioreanu" Oncology Institute, Bucharest, Romania

Rezumat

Fibroza retroperitoneală idiopatică (F.R.I) a preocupat lumea chirurgicală începând cu anul 1948 când a fost publicat, de către J. K. Ormond, primul caz de stenoză ureterală consecutivă unui proces sclero-inflamator retroperitoneal de cauză necunoscută. Până în 1960 s-au raportat în literatură 82 de cazuri. Autorii prezintă 2 noi cazuri personale de FRI, localizate exclusiv în compartimentul pelvin (subperitoneal) al spaţiului retroperitoneal. Este trecută in revistă literatura dedicată acestui subiect, se evocă teoriile etiologice actuale si se discută oportunităţile de tratament.

Cuvinte cheie: fibroza pelvină retroperitoneală, stenoză rectală